

ANNEX 5: Stock summary and assessment for the 2021 report

NAME	DEFINITION
TAXON_GROUP_NAME	Description of species
ICES_STOCK	ICES stock code
Opinion_type	1-A Analysis with Fmsy; 1-B Analysis without Fmsy; 2-A ICES advice without analysis; 2-B RFMO and GFCM advice without analysis; 3-A Expert assessment (with ICES advice).
Exploitation_level_compared_to	1 'exploited'; 2. 'not known'/'unclear'; 0 'overfished'
Stock_status	1 'healthy'; 2.0: 'not known'/'unclear'; 2.1 Status unknown but increasing; 2.2 Status unknown but decreasing; 2.3 Status unknown but stable; 0: 'in poor condition'.
Assessment_report_2020	1= 'exploited' or 'in good condition' 0= 'overfished' or 'in poor condition'. Stocks where F is poor but biomass is high where the B/'Bmsy' ratio is greater than 1.5 or very high where the ratio is close to or greater than 2, have been evaluated as being in good condition.
France's share of total landings of stock	France's share of total landings of stock. If figures were unavailable for international landings, the figures for France were used, thereby attributing 100% of landings to French fleet segments.
ICES_STOCK*	Stock for which France accounted for 100% only because in 2016 it was not possible to gather figures on total landings by other flag vessels.

Name	ICES stock	Opinion type	Level of exploitation	Stock status	Assessment for 2021 report	France's share of total landings of stock
Albacore	ALB (27)	1-A	1	1	1	22.6%
Alfonsino	ALF (27)	2-A	2	2	1	5.6%
European anchovy	ANE (VIII)	2-B	2	0	0	7.8%
	ANE (37.GSA7)	1-B	1	1	1	100%
Bigeye tuna	BET (47)	1-A	0	0	0	6.8%
	BET (51)	1-A	1	1	1	5.5%
Bluefin tuna	BFT (27)	1-A	1	1	1	5.2%
	BFT (37)	1-A	1	1	1	23.4%
Blue ling	BLI (I,II,IIIa,IVa,VIII,IX,XII)	2-A	2	0	0	7.5%
	BLI (Vb,VI,VII)	1-A	1	1	1	50.4%
Black scabbardfish	BSF (27)	2-A	1	2.3	1	28.6%
European seabass	BSS (IVbc,VIIa,VIIId-h)	1-A	1	0	1	35.1%
	BSS (VIIIab)	1-A	0	1	1	96.3%
Blue marlin	BUM (51)	1-A	0	1	0	1.9%

Atlantic cod	COD (IIIan,IV,VIIId)	1-A	0	0	0	1.4%
	COD (VIa)	1-A	0	0	0	7.1%
	COD (VIIe-k)	1-A	0	0	0	35.1%
Edible crab	CRE (VII,VIIIabd)	3-A	1	1	1	35.0%
European eel	ELE (27)	2-A	2	0	0	100.0%
	ELE (37)	2-A	2	0	0	51.7%
Greater forkbeard	GFB (27)	2-A	2	2.2	0	26.0%
Red gurnard	GUR (MI,IV,V,VI,VM,VMI)	2-A	2	2	1	61.5%
Haddock	HAD (VIIb-k)	1-A	0	1	1	59.5%
Atlantic herring	HER (IIIa,IV,VIIId)	1-A	1	1	1	5.5%
European hake	HKE (IMA,IV,VI,VM,VIMabd)	1-A	0	2.2	0	40.8%
	HKE (37.GSA7)	1-A	0	1	1	33.7%
Atlantic horse mackerel	HOM (Na,IVa,Vb,VIa,VNa-c,VMe-k,VI)	1-A	1	0	1	3.4%
Megrin	LEZ (IVa,VIa)	1-A	1	1	1	7.3%
	LEZ (VIIb-k,VIIIabd)	1-A	0	1	1	38.0%
Ling	LIN (IIIa,IVa,VI,VII,VIII,IX,XII,XIV)	2-A	1	2.1	1	10.9%
Atlantic mackerel	MAC (27)	1-A	0	0	0	2.1%
Monkfishes nei	MNZ (IIIa,IV,VI)	2-A	2	2.1	1	7.2%
	MNZ (VMb-k,VIMabd)	1-A	1	1	1	54.1%
	MNZ (VMb-k,VIMabd)	2-A	2	2	1	54.1%
Surmullet	MUR (VI,VIIa-c,VIIe-k,VIII,IXa)	1-A	0	1	0	73.2%
Red mullet	MUT (37.GSA7)	1-A	0	2.1	0	83.2%
Norway lobster	NEP (VIIbcjk.FU16)	1-A	1	0	1	2.4%
	NEP.fu20-21	1-A	0	2.3	1	6.5%
	NEP (VIIIab.FU23-24)	2-A	2	0	0	100%
Penaeus shrimp	PEN (31+41)	2-B	1	2.3	1	100%
European pilchard	PIL (VM,VNIabd)	1-A	0	1	0	97.2%
	PIL (37.GSA7)	1-A	1	1	1	59.2%
European plaice	PLE (VIIId)	2-A	0	1	1	47.4%
	PLE (VIIId)	2-A	0	0	0	6.2%
	PLE(7fg)	1-A	1	1	1	18.6%
	PLE (VIIhjk)	1-A	0	1	1	16.7%
Saithe	POK (IIIa,IV,VI)	2-A	1	2	1	13.3%

Pollack	POL (VI,VII)	2-A	2	2	1	25.3%
	POL (VIII,IXa)	2-A	2	2.1	1	79.0%
White skate	RJA (27)	2-A	2	2.1	1	75.1%
Thornback ray	RJC (INa,IV,VMd)	2-A	2	2	1	53.9%
	RJC (VI)	2-A	2	2.2	0	3.5%
	RJC (VIIafg)	2-A	2	2	1	12.2%
	RJC (VIIIe)	2-A	2	2	1	53.8%
	RJC (VIII)	2-A	2	2.1	1	43.4%
Shagreen ray	RJF (VI,VII)	2-A	2	2.1	1	58.9%
Sandy ray	RJI (VI,VII)	2-A	2	2.1	1	77.8%
Spotted ray	RJM (M1a,IV,VMd)	2-A	2	2.1	1	6.4%
	RJM (VI,VIIbj)	2-A	2	2.1	1	10.7%
	RJM (VNa,VNe-h)	2-A	2	2	1	81.6%
Cuckoo ray	RJN (VI,VM,VINabd)	2-A	2	2	1	71.2%
Undulate ray	RJU (VIIde)	2-A	1	1	1	74.1%
	RJU (VIIIab)	2-A	2	0	0	100%
Roundnose grenadier	RNG (I,II,IV,Va2,VIII,IX,XIVab2)	3-A	2	2.1	1	0.4%
	RNG (Vb,VI,VII,XMb)	3-A	1	1	1	20.7%
Blackspot seabream	SBR (VI,VII,VIII)	3-A	2	2	1	24.2%
Great Atlantic scallop	SCE (VIIId)	2-A	2	2.1	1	74.4%
	SCE (VIIe)	1-A	0	1	0	100%
	SCE (VIIIa)	2-A	2	2.1	1	100%
	SCE (VIIIb)	2-B	1	1	1	99.8%
Smooth hounds nei	SDV (27)	1-A	1	1	1	79.8%
Skipjack tuna	SKJ (34)	1-A	1	1	1	7.7%
	SKJ (51)	1-A	1	1	1	7.2%
Common sole	SOL (IV)	1-B	2	1	1	1.3%
	SOL (VIIId)	1-A	1	1	1	53.2%
	SOL (VIIe)	1-A	0	0	0	17.5%
	SOL (VIIIab)	1-A	1	1	0	89.3%
Swordfish	SWO (27)	2-A	2	2.1	1	0.8%
	SWO (37)	2-A	1	1	1	1.6%
	SWO (51)	3-A	2	2	1	2.7%
Tusk	USK (IIIa,IV,Vb,VIa,VII,VIII,IX,XIIb)	1-A	0	1	0	3.6%

Blue whiting	WHB (27)	1-A	0	1	0	1.1%
Whelk	WHE (VIIId)	2-A	2	2	1	59.1%
	WHE (VIIe)	1-A	1	0	1	80.9%
Whiting	WHG (IV,VIIId)	1-A	0	0	0	15.9%
	WHG (VIIbc,VIIe-k)	1-A	1.0.	0.2.	0	57.8%
	WHG (VIII,IXa)	2-A	3.1.	2.0.	1	87.2%
Yellowfin tuna	YFT (47)	1-A	1.0.	1.0.	1	13.4%
	YFT (51)	1-A	0.0.	0.0.	0	6.5%
Snappers	SNA (31)	1-A	1.0.	1.1.	1	100 %
Caribbean spiny lobster	SNA (31)	1-A	1.0.	1.1.	1	99.9%
Spotted spiny lobster	NLG (31)	1-A	1.0.	1.1.	1	99.8%