

NAME	DEFINITION
TAXON_GROUP_NAME	Description of species
ICES_STOCK	ICES stock code
Opinion_type	1-A: Analysis with Fmsy; 1-B: Analysis without Fmsy; 2-A: ICES advice without analysis; 2-B: RFMO and GFCM advice without analysis; 3-A: Expert assessment (with ICES advice).
Level_of_exploitation_	1: 'exploited'; 2: 'not known'/'unclear'; 0: 'overfished'
Stock_status	1: 'healthy'; 2.0: 'not known'/'unclear'; 2.1 Status unknown but increasing; 2.2 Status unknown but decreasing; 2.3 Status unknown but stable; 0: 'in poor condition'.
Assessment_for_2022_report	1= 'exploited' or 'in good condition' 0= 'overfished' or 'in poor condition'. Stocks where F is poor but biomass is high where the B/'Bmsy' ratio is greater than 1.5 or very high where the ratio is close to or greater than 2, have been evaluated as being in good condition.
France's share of total landings of stock	France's share of total landings of stock. If figures were unavailable for international landings, the figures for France were used, thereby attributing 100% of landings to French fleet segments.

Description	ICES stock	Opinion type	Level of exploitation	Stock status	Assessment for 2022 report	France's share of total landings of stock
Albacore	ALB (27)	1-A	1	1	1	15,2%
	ALB (51)	1-A	0	1	0	1,0%
European anchovy	ALF (27)	2-C	2	2	1	10,2%
	ANE (VIII)	1-B	1	1	1	0,2%
	ANE (37.GSA7)	1-B	1	2	1	100,0%
Rusty jobfish	ARQ (51.7)	1-A	1	1	1	100,0%
Crucifix sea catfish	AXP (31)	1-B	1	1	1	19,5%
Bigeye tuna	BET (47)	1-A	1	0	1	3,5%
	BET (51)	1-A	0	1	0	3,4%
Atlantic bluefin tuna	BFT (27)	1-A	1	2	1	1,5%
	BFT (37)	1-A	1	2	1	15,3%
Blue ling	BLI (I,II,IIIa,IVa,VIII,IX,XII)	3-A	2	0	0	7,5%
	BLI (Vb,VI,VII)	1-A	1	1	1	45,0%
Black scabbardfish	BSF (27)	3-A	1	2,3	1	21,8%
European seabass	BSS (IVbc,VIIa,VIIId-h)	1-A	1	0	1	45,5%
	BSS (VIIIab)	1-A	1	1	1	99,1%
Blue marlin	BUM (31)	1-A	0	0	0	18,2%
	BUM (51)	1-A	0	0	0	4,3%
Atlantic cod	COD (IIIan,IV,VIIId)	1-A	0	0	0	1,4%
	COD (VIa)	1-A	0	0	0	10,8%
	COD (VIIe-k)	1-A	0	0	0	40,3%
Edible crab	CRE (VII,VIIIabd)	3-A	0	0	0	35,4%
Brilliant pomfret	EBS (51.7)	1-A	1	1	1	100,0%
European eel	ELE (27)	3-A	2	0	0	100,0%
	ELE (37)	3-A	2	0	0	52,0%
Oblique-banded grouper	EZR (51.7)	1-A	1	1	1	100,0%
Greater forkbeard	GFB (27)	2-B	2	2,3	1	27,5%
Red gurnard	GUR (III,IV,V,VI,VII,VIII)	2-B	2	2,1	1	74,4%
Haddock	HAD (VIIb-k)	1-A	1	1	1	48,5%
Atlantic herring	HER (IIIa,IV,VIIId)	1-A	1	1	1	4,6%
European hake	HKE (IIIa,IV,VI,VII,VIIIabd)	1-A	1	1	1	41,1%
	HKE (VIIIc,IXa)	2-B	2	2,2	0	0,6%
	HKE (37.GSA7)	1-A	0	2	0	27,1%
Atlantic horse mackerel	HOM (IIa,IVa,Vb,VIa,VIIa-c,VIIe-k,VIII)	1-A	1	0	1	3,0%
Megrin	LEZ (IVa,VIa)	1-A	1	1	1	6,0%
	LEZ (VIIb-k,VIIIabd)	1-A	1	1	1	39,3%
Ling	LIN (IIIa,IVa,VI,VII,VIII,IX,XII,XIV)	2-B	2	2,1	1	12,1%
Goldbanded jobfish	LRI (51.7)	1-A	1	1	1	100,0%
Atlantic mackerel	MAC (27)	1-A	1	1	1	2,1%
Monkfishes nei	MNZ (IIIa,IV,VI)	2-B	2	2,2	0	7,6%
	MNZ (VIIb-k,VIIIabd)	1-A	1	1	1	52,4%
Surmullet	MUR (VI,VIIa-c,VIIe-k,VIII,IXa)	2-C	2	2	1	61,8%
Red mullet	MUT (37.GSA7)	1-A	0	1	0	80,7%

Norway lobster	NEP (VIIbcjk.FU16)	1-A	1	2	1	2,2%
	NEP.fu20-21	1-A	1	1	1	15,1%
	NEP (VIIIab.FU23-24)	1-A	1	2	1	100,0%
Spotted spiny lobster	NLG (31)	1-A	1	1	1	100,0%
Penaeus shrimp	PEN (31+41)	3-A	0	0	0	100,0%
European pilchard	PIL (VII,VIIIabd)	1-A	0	0	0	88,6%
	PIL (37.GSA7)	1-B	1	2	1	100,0%
European plaice	PLE (VIId)	1-A	1	1	1	50,2%
	PLE (VIIe)	2-A	0	1	1	6,1%
	PLE(7fg)	1-A	1	1	1	8,7%
	PLE (VIIhjk)	2-A	0	2,1	0	13,1%
Saithe	POK (IIIa,IV,VI)	1-A	0	0	1	14,4%
Pollack	POL (VI,VII)	3-A	1	2	1	26,2%
	POL (VIII,IXa)	2-C	2	2	1	76,6%
Thornback ray	RJC (IIIa,IV,VIId)	2-B	2	2,2	0	56,5%
	RJC (VI)	2-B	2	2,3	1	5,5%
	RJC (VIIafg)	2-B	2	2,3	1	8,8%
	RJC (VIIe)	2-C	2	2	1	56,9%
	RJC (VIII)	2-B	2	2,2	0	43,4%
Shagreen ray	RJF (VI,VII)	2-C	2	2	1	70,9%
Sandy ray	RJI (VI,VII)	2-C	2	2	1	76,2%
Spotted ray	RJM (IIIa,IV,VIId)	2-B	2	2,2	0	14,3%
	RJM (VI,VIIbj)	2-B	2	2,2	0	17,2%
	RJM (VIIa,VIIe-h)	2-B	2	2,3	1	77,7%
Cuckoo ray	RJN (VI,VII,VIIIabd)	2-B	2	2,1	1	80,8%
Undulate ray	RJU (VIId)	2-B	2	2,1	1	71,0%
	RJU (VIIIab)	2-C	2	2	1	98,3%
Roundnose grenadier	RNG (I,II,IV,Va2,VIII,IX,XIVab2)	2-C	2	2	1	1,7%
	RNG (Vb,VI,VII,XIIb)	2-C	2	2	1	32,4%
Blackspot seabream	SBR (VI,VII,VIII)	3-A	2	0	0	14,1%
Great Atlantic scallop	SCE (VIId)	3-A	1	1	1	79,3%
	SCE (VIIe)	3-A	1	1	1	100,0%
	SCE (VIIIa)	3-A	2	2	1	100,0%
	SCE (VIIIb)	3-A	2	2	1	98,8%
Smooth hounds nei	SDV (27)	2-B	2	2,2	0	82,0%
Indo-Pacific sailfish	SFA (51)	1-A	0	1	0	0,1%
Skipjack tuna	SKJ (34)	3-A	1	1	1	6,2%
	SKJ (51)	1-A	1	1	1	5,5%
Caribbean spiny lobster	SNA (31)	1-A	1	1	1	100,0%
Snapper	SNA (31)	1-A	1	1	1	100,0%
Common sole	SOL (VIId)	1-A	0	0	0	58,3%
	SOL (VIIe)	1-A	1	1	1	15,9%
	SOL (VIIIab)	1-A	0	0	1	93,2%
Swordfish	SWO (27)	1-A	1	1	1	0,8%
	SWO (37)	1-A	1	0	1	1,4%
	SWO (51)	1-A	1	1	1	3,5%
Tusk	USK (IIIa,IV,Vb,Vla,VII,VIII,IX,XIIb)	2-B	2	2,2	0	3,4%
Yellow-edge lyretail	VRL (51.7)	1-A	1	1	1	98,7%
Blue whiting	WHB (27)	1-A	0	1	1	0,9%
Whelk	WHE (VIId)	3-A	2	2	1	44,1%
	WHE (VIIe)	3-A	1	1	1	72,2%
Whiting	WHG (IV,VIId)	1-A	1	1	1	11,3%
	WHG (VIIbc,VIIe-k)	1-A	0	0	0	45,5%
	WHG (VIII,IXa)	2-A	1	2	1	97,9%
Yellowfin tuna	YFT (31)	1-A	1	1	1	4,3%
	YFT (47)	1-A	1	1	1	13,1%
	YFT (51)	1-A	0	0	0	5,8%
Acoupa weakfish	YNA (31)	2-B	2	2,3	1	22,9%
Green weakfish	YNV (31)	1-B	1	1	1	16,0%